

Aspects of the Spin Dynamics in the Cuprate Superconductors

H. Monien

Institute for Theoretical Physics

University of California

Santa Barbara, CA 93106, USA

February 1, 2008

Abstract:

One of the interesting aspects of the CuO superconductors is that superconductivity is happening so close to the antiferromagnetic state. The nuclear magnetic resonance and the recent neutron scattering experiments clearly indicate that magnetic correlations persist in to the heavily doped regime. In this paper we will discuss some of the details of the coupling of the nuclear magnetic spin to the conduction electron spins. Furthermore we will show that a simple band structure can explain the recent neutron scattering data in the $\text{LaSr}_x\text{Cu}_{2-x}\text{O}_4$ material for the optimal concentration of $x \approx 0.15$ if the lifetime effects are included.

1 Introduction

From the theoretical perspective one of the most interesting aspects of the CuO high temperature superconductors is that they belong to a class of materials which show strong electronic correlations. In the undoped materials these correlations lead to the antiferromagnetic ordering of the electronic spins. This is why Anderson [1] pointed out early on that the physics of the CuO superconductors is best described in terms of the Hubbard model. The model Hamiltonian contains the strong on site Coulomb repulsion and a hopping term describing the kinetic energy. The theoretically fascinating aspect is how to obtain superconductivity from a purely repulsive interaction. Very close to half filling where this model describes the insulating Heisenberg antiferromagnet it is extremely well understood and the neutron scattering experiments [2] and theoretical calculations, e.g. [3], are in beautiful agreement. Away from half filling the situation is much less clear. Except in one dimensions and recently infinite dimensions the physics of the Hubbard model is not understood to a degree that one knows the correlation functions quantitatively. Even today small cluster calculations and approximations which are uncontrolled in the relevant regime are the only way one is able to achieve some progress in the strongly correlated regime away from half filling.

Under these circumstances it is useful to know what limitations one can obtain from experiments for a theory for the CuO superconductors. The constraints obtained from experiments might tell us if a theoretical model like the Hubbard model contains even qualitatively the right physics to describe the low energy excitations of the “real materials” or if some additional physics is required to understand the basic properties of the high temperature superconductors. It is quite clear that these materials are close to many instabilities which might all be important. Thus, choosing to concentrate on a particular degree of freedom of the system, one might completely miss the relevant physics. A “worst case scenario” from a theorists point of view would be that all the possible degrees of freedom of the system contribute to the most interesting of all the instabilities of the system: the superconducting state.

In order to see “a way through the jungle”, I will concentrate on the magnetic properties of the copper oxide superconductors which are very interesting in themselves. The magnetic properties of the CuO materials were studied in great detail by the nuclear magnetic resonance techniques by many groups and by neutron scattering experiments which are very difficult to perform in these compounds in comparison with standard materials. In both types of experiments one would like to obtain information about the dynamical structure factor of the electronic spins. Whereas the nuclear magnetic resonance experiments probe the local environment of one particular nucleus and therefore a wavevector average of the dynamical structure factor at very low energy transfers, the neutron scattering experiments can, at least in principle, scan all frequencies and wavevectors corresponding to the whole Brillouin zone.

The interpretation of the nuclear magnetic resonance experiments is quite in-

volved. It requires the knowledge of the so called hyperfine Hamiltonian which describes the coupling of the electronic spin to the nuclear spin. By now it is established that the spin degrees of freedom in the CuO superconductors can be described in a so called “one component picture”, which means that it is sufficient to consider one spin degree of freedom per unit cell. If one assumes a band structure picture one can say more precisely that only one band plays a role for the low energy physics of the spins. The nature of the band, i.e. the percentage of copper or oxygen admixture might vary, as one moves along the Fermi surface. But the statement about one component is more general in that it does not require a Fermi liquid state. The spin degree of freedom might equally well be carried by the strongly correlated quasiparticles of a tJ - model. The question is how do the spin of the quasiparticle couple to the nuclear spins. This is discussed in the next section.

2 Transferred hyperfine interactions and band structure

In this section we discuss the derivation of a hyperfine Hamiltonian similar to the so called Mila-Rice Hamiltonian [4] which was originally introduced to understand the difference in the anisotropies of the Cu nuclear magnetic relaxation rates and the Cu Knight shift measurements in terms of uncorrelated spins. The importance of a transferred hyperfine interaction in the case where the electronic spins are strongly correlated was pointed out early on by Shastry [5]. In his paper he showed that within the theoretical framework of the so called t-J model one can understand the difference between the magnitude of the Cu and O relaxation rates if the oxygen nuclear spins are only coupled to the electronic spins via a transferred hyperfine coupling.

The phenomenological hyperfine Hamiltonian which is compatible with this assumption of a “one component picture” is:

$$H_{hf} = \sum_{i\alpha\beta} {}^{63}I_{i\alpha\beta} A_{\alpha\beta} S_{i\beta} + B \sum_{\langle ij \rangle > \alpha} {}^{63}I_{i\alpha} S_{j\alpha} + \sum_{\langle ij \rangle > \alpha\beta} {}^{17}I_{i\alpha} C_{\alpha\beta} S_{j\beta} \quad (1)$$

where ${}^{63}I_{i\alpha}$ is the α th, $\alpha \in \{x, y, z\}$, component of the copper nuclear spin at the site i , $S_{i\alpha}$ is the α th component of the electronic spin, $A_{\alpha\beta}$ is the anisotropic copper hyperfine coupling, B is the transferred Cu hyperfine coupling and $C_{\alpha\beta}$ is the transferred oxygen hyperfine coupling.

This is the so called Mila-Rice Hamiltonian. The unusual feature is the large so called transferred hyperfine coupling B . Couplings to other nuclei can be included in a similar way. Whereas the Mila-Rice Hamiltonian was derived from a quantum-chemistry calculation on a small cluster and therefore in a very ionic picture Shastry assumed an itinerant electron system. It was argued that the success of the Mila-Rice Hamiltonian indicates that the Cu spins are nearly localized. Here I would like to point out that in an itinerant “band structure picture” one basically obtains the

same results for the hyperfine Hamiltonian. These results are mostly due to Takigawa [6]. In addition to the Mila Rice Hamiltonian one finds additional terms which are of similar size as the nearest neighbor transferred hyperfine interaction but we will argue that these terms are negligible because the correlations are only short ranged.

The starting point of our calculation is the canonical band structure for the plane. We include in addition to copper $3d_{x^2-y^2}$, the oxygen $2p_x$ and $2p_y$ orbitals and the copper $4s$ orbital. Here it is sufficient to include the $4s$ orbital only in perturbation theory. The Hamiltonian has the form

$$H_0 = \sum_{k\alpha} \{ \epsilon_d d_{k\alpha}^\dagger d_{k\alpha} + \epsilon_p p_{x,k\alpha}^\dagger p_{x,k\alpha} + \epsilon_p p_{y,k\alpha}^\dagger p_{y,k\alpha} \} + \sum_{k\alpha} 2it d_{k\alpha}^\dagger (\sin(k_x/2) p_{x,k\alpha} + \sin(k_y/2) p_{y,k\alpha}) + h.c. \quad (2)$$

where $d_{k\alpha}^\dagger$ ($d_{k\alpha}$) is the copper $3d_{x^2-y^2}$ electron creation (annihilation) operator, ϵ_d is the energy of that orbital, $p_{x,y,k\alpha}^\dagger$ are the annihilation operators for the oxygen (p_x , p_y) orbitals in the plane. If the $4s$ orbital is partially filled the electron spin has a contact interaction with the nuclear spin at site i which is given by:

$$H_{hf} = B \cdot {}^{63}\vec{I}_i \cdot s_{i\alpha}^\dagger \vec{\sigma}_{\alpha\beta} s_{i\beta} \quad (3)$$

where $s_{i\alpha}^\dagger$ are the annihilation (creation) operators for the $4s$ orbital at the Cu site i , $\vec{\sigma}$ is the vector of Pauli matrices and ${}^{63}\vec{I}_i$ is the Cu nuclear spin. The coupling constant B depends on atomic physics and can be found for example in [4]. The contact interaction is isotropic and therefore the transferred hyperfine coupling will be isotropic. If we believe that only one band of the three bands of our model plays a role for the low energy physics we would like to reduce hyperfine Hamiltonian to the part which is connected with this one band. Therefore we project the hyperfine Hamiltonian, Eq. 3, on to the conduction electron band:

$$(H_{hf})_{proj} = \sum_{kk'} \langle k\alpha | H_{hf} | k'\alpha' \rangle c_{k\alpha}^\dagger c_{k'\alpha'} \quad (4)$$

where $c_{k\alpha}^\dagger$ are the creation and annihilation operators of the conduction electrons with momentum $\vec{k}$ and spin α .

The projected hyperfine Hamiltonian has the form:

$$(H_{hf})_{proj} = B \cdot {}^{63}\vec{I}_0 \cdot \sum_{ij} f_i^* f_j c_{i\alpha}^\dagger \vec{\sigma}_{\alpha\beta} c_{j\beta} \quad (5)$$

where f_i is the Fourier transform of $f_k = \langle 4s | k \rangle$, the overlap between the $4s$ state and the conduction electron bands at the site i . It turns out that we only need to know this overlap. This is basically what Mila and Rice did for their ionic model. The parameters for a fit of tight binding model to the “real band structure” were given by e.g. Hybertsen and Schlüter [7]. For the energy of the Cu d orbital, the oxygen

p orbital and the Cu $4s$ orbital respectively they found $\epsilon_d = -2.1\text{eV}$, $\epsilon_p = -3.3\text{eV}$ and $\epsilon_{4s} = -8.1\text{eV}$. For the overlaps they give the following numbers $t_{pd} = 1.3\text{eV}$, $t_{pp} = 1\text{eV}$ and $t_{ps} = 3.8\text{eV}$. With this parameters we can calculate the hyperfine interactions (for details consult e.g. [8]). In addition to the terms which couple the conduction electron spin on one site to the nuclear spin on a neighboring site we find terms which describe a nuclear spin assisted spin flip + hop term. The result of the calculation can be cast into the simple form:

$$(H_{hf})_{proj} = B_{10,10} {}^{63}\vec{I}_{00} \cdot \vec{S}_{11} + B_{10,01} {}^{63}\vec{I}_{00} \cdot c_{10,\alpha}^\dagger \vec{\sigma}_{\alpha\beta} c_{01,\beta} + \dots \quad (6)$$

where the coupling constants are of the order $B_{10,10} \sim 48\text{kOe}/\mu_B$, $B_{10,01} = B_{10,10}$. The coupling constant $B_{10,10}$ is the transferred hyperfine coupling for the Cu nuclear spin as discussed by Mila and Rice, the spin flip hop term $B_{10,01}$ is not present in the ionic picture. The next term in the expansion is about a factor of 6 smaller than the first term and can be neglected safely. Although the nuclear spin assisted spin flip hop term is of the same magnitude as the transferred term we can neglect it in our analysis of the nuclear magnetic resonance experiments because the dynamical structure factor does not have large weight at this particular area of momentum space. Therefore we are left with the standard hyperfine Hamiltonian for the CuO plane.

If we do not allow for any other relaxation channel which might for example be produced by a coupling of the nuclear spin to the orbital motion of the electron then we know exactly which moments of the dynamical structure factor contribute to the nuclear magnetic relaxation.

3 How important is orbital relaxation?

In previous theories of the nuclear magnetic relaxation in the high temperature superconductors it was assumed the only way the nuclear spin couples to the electrons is via the electronic spin. The unusual behavior of the anisotropy of the Cu nuclear magnetic relaxation rate as a function of temperature [9], [10] raises the question if there is another relaxation channel present. In a recent paper Millis and Monien [11] discuss the possibility of another relaxation channel for the Cu nuclear spin, namely the coupling of the Cu nuclear spin to the orbital motion of the electron which was already mentioned by Warren and Walstedt in an early paper. The coupling of the nuclear spin to the orbital momentum is given by:

$$H_{\text{Orb}} = {}^{63}\gamma_n \gamma_e \hbar^2 \frac{\vec{I} \cdot \vec{L}}{r^3} \quad (7)$$

where $\vec{I}$ is the copper nuclear spin, $\vec{L}$ the electron angular momentum and ${}^{63}\gamma_n$ and γ_e the gyromagnetic ratio of the copper nucleus and the electron respectively. In order to estimate the size of this effect we use again the simplest tight binding picture for the CuO band of a single plane with the same parameter as used for the calculation of the

hyperfine Hamiltonian. For the calculation of the relaxation rates we need to know the matrix elements of the orbital coupling, Eq. (7), with the Wannier functions. The rates can then be calculated simply by using Fermis “golden rule”. At a temperature of about 100K we find for the nuclear magnetic relaxation rate produced by *orbital currents* for a field applied parallel and perpendicular to the CuO plane:

$$\begin{aligned} W_{\parallel} &= 3.8 \times C_{x^2-y^2} (4C_{xy} + C_{xz}) \text{ ms}^{-1} \\ W_{\perp} &= 3.8 \times C_{x^2-y^2} C_{xz} \text{ ms}^{-1} \end{aligned} \quad (8)$$

The coefficients $C_{x^2-y^2}$, C_{xy} and C_{xz} denote the percentage of admixture of the Cu $3d_{x^2-y^2}$, $3d_{xy}$ and $3d_{xz}$ orbital respectively. The result of the analysis is that the orbital relaxation of the nuclear spins can be appreciable because of the orbital degeneracy but that the absolute value depends strongly on the exact amount of admixture of the Cu $3d_{xz}$ and $3d_{xy}$ orbitals to the Wannier functions at the Fermi surface which are not known theoretically to an accuracy, ($\sim 5\%$), needed to determine if the orbital relaxation may play a role or not. Nevertheless it is important to keep this additional relaxation channel in mind if one discusses the limitations on hyperfine Hamiltonian and dynamical structure factor obtained from experiments.

4 Reexamination of the MMP model

Up to now we have only discussed the coupling of the electronic spin and orbital degree of freedom to the nuclear spin. Now we would like to turn our attention to the quantity which is of most interest from the theoretical point of view, the dynamical structure factor.

Recently we [8] reexamined the results of a previous phenomenological analysis of the nuclear magnetic relaxation experiments in the $\text{YBa}_2\text{Cu}_3\text{O}_{6+\delta}$ materials [12]. In the original MMP paper we made the assumption that the only temperature dependent quantity is the antiferromagnetic correlation length and the energy scale is described by a dynamical critical exponent and used the simplest form for a peaked structure factor describing the magnetic correlations, a Lorentzian. In the extended analysis we dropped the assumption that the correlation length is the only quantity which is temperature dependent and allowed the strength of the antiferromagnetic peak also to vary with temperature. Also we considered two models for the q dependence of the structure factor - a Lorentzian and a Gaussian. The parameters describing the correlation functions are the antiferromagnetic correlation length ξ and the strength of the antiferromagnetic peak which we denote with β . The two models for the dynamical structure factor in the limit frequency ω going to zero are:

$$\chi''_{Lor}(q, \omega) = \frac{\pi\chi_0\hbar\omega}{\Gamma} \left[1 + \beta \frac{\xi^4}{(1 + \xi^2(q - Q)^2)^2} \right] \quad (9)$$

$$\chi''_{Gauss}(q, \omega) = \frac{\pi\chi_0\hbar\omega}{\Gamma} \left[1 + \beta\xi^4 e^{-\xi^2(q-Q)^2} \right] \quad (10)$$

where ξ is the antiferromagnetic correlation length, β is the strength of the antiferromagnetic peak at the zone corner, $Q = (\pi/a, \pi/a)$, Γ is the “bare” energy scale of the spin fluctuations and χ_0 is the static susceptibility.

Within the “one component picture” the Knight shift for a field applied in the direction α , K_α , and the relaxation rates, W_α of the various nuclei, denoted by $a = 63, 17, 89$ for Cu, O and Y respectively, for a field applied in the α direction are given by:

$$\begin{aligned} {}^a K_\alpha &= \lim_{q \rightarrow 0} {}^a F_\alpha(q) \chi'(q, \omega = 0) \\ {}^a W_\alpha &= \frac{1}{4\mu_B^2} \lim_{\omega \rightarrow 0} \frac{k_B T}{\omega} \sum_{q, \bar{\alpha}} [{}^a F_{\bar{\alpha}}]^2 \chi''(q, \omega) \end{aligned} \quad (11)$$

where $\bar{\alpha}$ is the direction perpendicular to α . The form factors ${}^a F_\alpha(q)$ have the dimension of energy and are basically the Fourier transform of the spin hyperfine Hamiltonian and can be found in the literature, e.g. [8]. To obtain the limits on the correlation lengths we considered two limits in one case the correlation length is temperature dependent and the strength is not. In the second case the strength of the peak is temperature dependent and the correlation length is temperature independent.

We state the results for the correlation lengths in Table (1).

	YBa ₂ Cu ₃ O _{6.63}		YBa ₂ Cu ₃ O ₇	
T[K]	ξ_{Lor}	ξ_{Gauss}	ξ_{Lor}	ξ_{Gauss}
100	1.5 - 2.4	1.0 - 1.5	2.5 - 4.5	1.5 - 2.5
300	1.3 - 1.7	0.7 - 1.1	1.5 - 2.5	1.0 - 1.5

Table 1: correlation lengths from the NMR analysis

From these results we can conclude:

- The results for the correlation length are not compatible with the neutron scattering results by Rossat - Mignod [13].
- The neutron scattering experiments favor temperature independent correlation lengths.
- The experimental results by G. Aeppli and coworkers [14] seem to be more compatible with the correlation lengths required by NMR experiments.

The detailed analysis shows that the copper and oxygen relaxation time experiments are compatible with both assumptions - a temperature dependent correlation length or a temperature dependent strength of the antiferromagnetic peak - but that the crucial quantity which could determine which case it is is the ratio of the yttrium

to the oxygen relaxation rate which was not measured in the same sample to a high enough accuracy up to now. For details of the analysis we refer the reader to our paper [8]. The original MMP analysis was intended to describe the spin correlations on a very low energy scale. A comparison of the nuclear magnetic experiments with the neutron scattering experiments requires an extrapolation to larger energies. For a thorough discussion we refer the reader to [8]

5 Dynamical structure factor for in the marginal Fermi liquid picture

The remaining physics question, as stated in the introduction, is off course where does one get the low energy physics scale from. The easiest way to obtain a low energy scale from a “high energy” model is to be close to a critical point in this class of models belong the “nested Fermi liquid”, Hubbard RPA and dynamical phase separation models. Very close to the phase transition dynamical scaling should hold in that case the low energy scale is determined by the correlation length via $\omega \sim \Gamma^{-z}$ where Γ is the energy scale determined by the “high energy physics” and z is the dynamical scaling exponent. In this section we will circumnavigate the difficult question what is determining the low energy physics and answer the following related question: Assuming that the only effect of all the many body complications is to give rise to a single particle scattering rate which is proportional to the energy or temperature, whatever is less (which is the so called marginal Fermi liquid hypothesis) what can we know about the spin response. The original marginal Fermi liquid picture, [15], assumed that the selfenergy is nearly wavevector independent

$$\text{Im}\Sigma \sim \max(\omega, T). \quad (12)$$

In the original MFL paper it was argued that the resulting response functions are a universal function of ω/T where ω is the frequency of the external field and T is the temperature and also weakly q dependent. The assumed factorizable form of the dynamical structure factor is not consistent with a different temperature dependence of the Cu and O relaxation rate. Neutron scattering experiments by Aeppli et al. [14] on the $\text{LaSr}_x\text{Cu}_{2-x}\text{O}_4$ for $x \approx 0.15$ showed that the dynamical structure factor is strongly q dependent even for the strongly doped material. Littlewood et al., [16] have proposed an extension of the original MFL picture. It is clear that in two dimensions the dynamical structure factor even for the noninteracting Fermi gas shows a strong q dependence at low frequencies. It is interesting to examine the predictions of the simple band structure of the CuO plane. Using a tight binding band structure one obtains an incommensurate peak structure with four posts which are a little bit shifted away from the zone corner, $Q = (\pi/a, \pi/a)$. What we found is that the position and magnitude seems to be compatible with a simple tight binding band structure picture for the CuO plane. The position and magnitude of the peaks

will strongly depend on the doping. What is even more interesting is that if we include the marginal Fermi liquid form for the self energy we also can understand the temperature and energy dependence of the peak structure. In the regime which is accessible by the neutron scattering the main effect of the marginal Fermi liquid form of the quasiparticle propagator is the life time broadening which has the energy scale of the temperature.

Figure 1: The dynamical structure factor $S(q, \omega)$ for various energy transfers, $\omega = 3, 6, 12$ and 15 meV as a function of the momentum transfer in the q_x direction. q_y is fixed to π/a . The bottom curve corresponds to the lowest energy transfer the top to the highest.

The magnetic properties of the heavily doped regime of $\text{LaSr}_x\text{Cu}_{2-x}\text{O}_4$ can therefore be understood in terms of a simple band structure picture of marginal Fermi liquid quasiparticles.

6 Conclusions

By now it has become quite clear that temperature dependent antiferromagnetic correlations play an important role in the understanding of the magnetic properties of the CuO high temperature superconductors even in the strongly doped regime. The early interpretation of Millis, Monien and Pines [12] assumed a temperature dependent correlation length. The reexamination of the present nuclear magnetic resonance data by [8] demonstrates that one can not deduce from the nuclear magnetic relaxation measurements of the copper and oxygen rates alone that the correlation length is temperature dependent. We have proposed a simple model in which the temperature and frequency dependence of the dynamical structure factor in $\text{LaSr}_x\text{Cu}_{2-x}\text{O}_4$, $x \approx 0.15$,

can be understood in a simple tight binding band structure picture including lifetime effects.

It is a pleasure to thank my numerous collaborators. Many have worked in this interesting area of solid state physics - I apologize to those whose important work I did omit in this brief review. The research for this project was supported in part by an IBM fellowship and by the NSF Grant NSF-PHY-04035.

References

- [1] P. W. Anderson, Science **235**, 1196 (1989)
- [2] see for e.g. B. Keimer et al., Phys. Rev. Lett. **67**, 1930 (1991)
- [3] For a review see, e.g. S. Chakravarty, "Spin Dynamics of Superconducting Cuprates", p. 198 in High Temperature Superconductivity: Proceedings of the Los Alamos Symposium - 1989, K. S. Bedell, D. Coffey, D. E. Meltzer, D. Pines and J. R. Schrieffer, eds. (Addison Wesley: Redwood City, CA) (1990).
- [4] F. Mila and T. M. Rice, Physica **C 157**, 561 (1989)
- [5] B. S. Shastry, Phys. Rev. Lett. **63**, 1288 (1989)
- [6] M. Takigawa, private communication
- [7] see e.g. M. Schlüter and M. S. Hybertsen, Physica **C 162**, 583 (1989)
- [8] A. J. Millis and H. Monien, Phys. Rev. B **B45**, 3059 (1992)
- [9] M. Takigawa, J. L. Smith and W. L. Hults, Phys. Rev. **B44**, 7764 (1991)
- [10] J. A. Martindale et al., Phys. Rev. Lett. **68**, 702 (1992)
- [11] A. J. Millis and H. Monien, unpublished
- [12] A. J. Millis, H. Monien and D. Pines, Phys. Rev. **B42**, 167 (1990)
- [13] see e.g. J. Rossat-Mignod et al., Physica **C185**, 86 (1991)
- [14] T. E. Mason, G. Aeppli, M.A. Mook, Phys. Rev. Lett. **68**, 1414 (1992)
- [15] C. M. Varma et al., Phys. Rev. Lett. **63**, 1996 (1989)
- [16] P. Littlewood, J. Zaanen, G. Aeppli and H. Monien, unpublished